

Breathing Signature of the Core Promoter

B. S. Alexandrov¹, V. Gelev², A. R. Bishop¹, K. Ø. Rasmussen¹, and A. Usheva²

We report our finding that the introduction of a few mismatched bases (artificial “bubble template”) at the transcription start site allows bidirectional transcription even in the absence of any transcription factors. We establish the correlation between experimentally determined DNA gene transcription regulation sites and the theoretically determined thermally induced coherent openings (bubbles) of the double-stranded DNA at these sites. Our analysis suggests a complex dynamic picture of DNA promoter regions, where different regulatory sites may be distinguishable by the details of their dynamic behavior.

Keywords — DNA, promoters, local dynamics, bubbles, transcription.

I. TRANSCRIPTION BUBBLE

From a structural perspective, RNA polymerase require single-stranded DNA, i.e. the formation of a ‘transcriptional bubble’ at the transcriptional start site (TSS) to initiate transcription [1]. Eukaryotic transcription initiation often proceeds from a negatively supercoiled template in the absence of helicases [2–5], suggesting spontaneous local melting of double-stranded (ds)DNA as a key feature of promoter sequences. Furthermore, the introduction of a few mismatched bases to permanently unzip the DNA at the start site allows transcription in the absence of supercoiling [5,6].

We attempt to better establish the connection between DNA bubbles and transcription through the following simulations and experiments.

II. SIMULATIONS

Since localized breathing of the double strand is the dynamical mechanism of bubble generation it is reasonable to expect some local enhancement of the breathing dynamics to be a common feature of the TSS, in order to assist the formation of the transcriptional bubble. Previously we have demonstrated a correlation between TSS location, single strand nuclease sensitivity, and transient dsDNA strand separation [7], as well as correlation between enhanced bubble formation and various transcription binding sites [8]. In search of distinguishing dynamic features of gene promoter TSS sequences, we performed molecular dynamic simulations, based on the Peyrard-Bishop-Dauxois model [9,10] of eight experimentally characterized mammalian core promoters [11]. From the generated dynamical trajectories we extracted three distinct dynamic characteristics: bubble probability, bubble lifetime, and the average strand separation. The calculated dynamical profiles

suggest that thermal fluctuations commonly induce relatively large, long-lived bubbles at the transcription start site [11].

III. EXPERIMENTS

Experimentally, we demonstrated that human RNA polymerase (RNAP) II performs bidirectionally transcription in the absence of any transcription factors, if an artificial long-lived bubble of length more than 5 base pairs is introduced at the start site of the Adeno-associated virus P5 promoter [11]. This transcriptional data together with the previously published results by us and others [2-7], clearly suggests that most likely local DNA melting is sufficient to enable bidirectional transcription by RNAP II alone.

IV. CONCLUSIONS

Our data, suggest that each promoter exhibits distinct DNA dynamic characteristics, i.e. spontaneous local openings, which are likely to be recognized and engaged by the transcriptional machinery, and may then be amplified, stabilized, or suppressed by DNA-protein interactions as part of gene transcriptional regulation.

REFERENCES

- [1] Cheetham GM, Zeruzalmi D, Steitz TA (1999) *Structural basis for initiation of transcription from an RNA polymerase-promoter complex*. Nature 399: 80–83.
- [2] Timmers HT (1994) *Transcription initiation by RNA polymerase II does not require hydrolysis of the beta-gamma phosphoanhydride bond of ATP*. EMBO J 13: 391–399.
- [3] Parvin JD, Sharp PA (1993) *DNA topology and a minimal set of basal factors for transcription by RNA polymerase II*. Cell 73: 533–540.
- [4] Leblanc Bp Fau, Benham Cj Fau, Clark DJ (2000) *An initiation element in the yeast CUP1 promoter is recognized by RNA polymerase II in the absence of TATA box-binding protein if the DNA is negatively supercoiled*. Proc Natl Acad Sci U S A 97: 10745–10750.
- [5] Usheva A, Shenk T (1996) *YY1 transcriptional initiator: protein interactions and association with a DNA site containing unpaired strands*. Proc Natl Acad Sci U S A 93: 13571–13576.
- [6] Pan G, Greenblatt J (1994) *Initiation of transcription by RNA polymerase II is limited by melting of the promoter DNA in the region immediately upstream of the initiation site*. J Biol Chem 269: 30101–30104.
- [7] Choi CH, Kalosakas G, Rasmussen KO, Hiromura M, Bishop AR (2004) *DNA dynamically directs its own transcription initiation*. Nucleic Acids Res 32: 1584–1590.
- [8] Choi CH, Rapti Z, Gelev G, Hacker MR, Alexandrov BS, Park EJ, Park JS, Horikoshi N, Smerzi A, Rasmussen KO, Bishop AR, Usheva A (2008) *Profiling the Thermodynamic Softness of Adenoviral Promoters*, Biophys. J 95, 597.
- [9] Peyrard M, Bishop AR (1989) *Statistical mechanics of a nonlinear model for DNA denaturation*. Phys Rev Lett 62: 2755–2758. Dauxois T, Peyrard M, Bishop AR (1993) *Entropy-driven DNA denaturation*. Phys Rev E 47: R44.
- [10] Alexandrov BS, Wille LT, Rasmussen KO, Bishop AR, Blagoev KB (2006) *Bubble statistics and dynamics in double-stranded DNA*. Phys Rev E 74: 050901.
- [11] Alexandrov BS, Gelev V, Wook Yoo S, Bishop AR, Rasmussen KO, Usheva A, (2009) *Toward a Detailed Description of the Thermally Induced Dynamics of the Core Promoter*, PLoS Computational Biology, 5:e1000313.

This work was funded by NIH grant GM071482, and supported by DOE

¹ Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico, United States of America,

² Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, United States of America

